

# A note on random minimum length spanning trees

Alan Frieze\*      Miklós Ruzinkó†

Lubos Thoma‡

Department of Mathematical Sciences

Carnegie Mellon University

Pittsburgh PA15213, USA

June 27, 2000

## Abstract

Consider a connected  $r$ -regular  $n$ -vertex graph  $G$  with random independent edge lengths, each uniformly distributed on  $[0, 1]$ . Let  $mst(G)$  be the expected length of a minimum spanning tree. We show in this paper that if  $G$  is sufficiently highly edge connected then the expected length of a minimum spanning tree is  $\sim \frac{n}{r}\zeta(3)$ . If we omit the edge connectivity condition, then it is at most  $\sim \frac{n}{r}(\zeta(3) + 1)$ .

## 1 Introduction

Given a connected simple graph  $G = (V, E)$  with edge lengths  $\mathbf{x} = (x_e : e \in E)$ , let  $mst(G, \mathbf{x})$  denote the minimum length of a spanning tree. When  $\mathbf{X} = (X_e : e \in E)$  is a family of independent random variables, each uniformly distributed on the interval  $[0, 1]$ , denote the expected value  $\mathbf{E}(mst(G, \mathbf{X}))$  by  $mst(G)$ . Consider the complete graph  $K_n$ . It is known (see [2]) that, as  $n \rightarrow \infty$ ,  $mst(K_n) \rightarrow \zeta(3)$ . Here  $\zeta(3) = \sum_{j=1}^{\infty} j^{-3} \sim 1.202$ . Beveridge, Frieze and McDiarmid [1] proved two theorems that together generalise the previous results of [2], [3], [5].

**Theorem 1** *For any  $n$ -vertex connected graph  $G$ ,*

$$mst(G) \geq \frac{n}{\Delta}(\zeta(3) - \epsilon_1)$$

where  $\Delta = \Delta(G)$  denotes the maximum degree in  $G$  and  $\epsilon_1 = \epsilon_1(\Delta) \rightarrow 0$  as  $\Delta \rightarrow \infty$ .

---

\*Supported in part by NSF Grant CCR9818411 email: alan@random.math.cmu.edu

†Permanent Address Computer and Automation Research Institute of the Hungarian Academy of Sciences, Budapest, P.O.Box 63, Hungary-1518. Supported in part by OTKA Grants T 030059 and T 29074 FKFP 0607/1999. email: ruzinko@lutra.sztaki.hu

‡Supported in part by NSF grant DMS-9970622. email: thoma@qwes.math.cmu.edu

For an upper bound we need expansion properties of  $G$ .

**Theorem 2** *Let  $\alpha = \alpha(r) = O(r^{-1/3})$  and let  $\rho = \rho(r)$  and  $\omega = \omega(r)$  tend to infinity with  $r$ . Suppose that the graph  $G = (V, E)$  is connected and satisfies*

$$r \leq \delta \leq \Delta \leq (1 + \alpha)r, \quad (1)$$

where  $\delta = \delta(G)$  denotes the minimum degree in  $G$ . Suppose also that

$$|(S : \bar{S})|/|S| \geq \omega r^{2/3} \log r \text{ for all } S \subseteq V \text{ with } r/2 < |S| \leq \min\{\rho r, |V|/2\}, \quad (2)$$

where  $(S : \bar{S}) = \{(x, y) \in E : x \in S, y \in \bar{S} = E \setminus S\}$ . Then

$$\left| mst(G) - \frac{n}{r} \zeta(3) \right| \leq \epsilon_2 \frac{n}{r}$$

where the  $\epsilon_2 = \epsilon_2(r) \rightarrow 0$  as  $r \rightarrow \infty$ .

For regular graphs we of course take  $\alpha = 0$ .

The expansion condition in the above theorem is probably not the “right one” for obtaining  $mst(G) \sim \frac{n}{r} \zeta(3)$ . We conjecture that high edge connectivity is sufficient: Let  $\lambda = \lambda(G)$  denote the *edge connectivity* of  $G$ .

**Conjecture 1**

*Suppose that (1) holds. Then,*

$$\left| mst(G) - \frac{n}{r} \zeta(3) \right| \leq \epsilon_3 \frac{n}{r}$$

where  $\epsilon_3 = \epsilon_3(\lambda) \rightarrow 0$  as  $\lambda \rightarrow \infty$ .

Note that  $\lambda \rightarrow \infty$  implies  $r \rightarrow \infty$ .

Along these lines, we prove the following theorem.

**Theorem 3** *Assume  $\alpha = \alpha(r) = O(r^{-1/3})$  and (1) is satisfied. Suppose that  $r \geq \lambda(G) \geq \omega r^{2/3} \log n$  where  $\omega = \omega(r)$  tends to infinity with  $r$ . Then*

$$\left| mst(G) - \frac{n}{r} \zeta(3) \right| \leq \epsilon_4 \frac{n}{r}$$

where the  $\epsilon_4 = \epsilon_4(r) \rightarrow 0$  as  $r \rightarrow \infty$ .

**Remark:** It is worth pointing out that it is not enough to have  $r \rightarrow \infty$  in order to have the result of Theorem 2, that is, we need some extra condition such as high edge connectivity. For consider the graph  $\Gamma(n, r)$  obtained from  $n/r$   $r$ -cliques  $C_1, C_2, \dots, C_{n/r}$  by deleting an edge  $(x_i, y_i)$  from  $C_i$ ,  $1 \leq i \leq n/r$  then joining the cliques into a cycle of cliques by adding edges  $(y_i, x_{i+1})$  for  $1 \leq i \leq n/r$ . It is not hard to see that

$$mst(\Gamma(n, r)) \sim \frac{n}{r} \left( \zeta(3) + \frac{1}{2} \right)$$

if  $r \rightarrow \infty$  with  $r = o(n)$ . We repeat the conjecture from [1] that this is the worst-case, i.e.

**Conjecture 2** *Assuming only the conditions of Theorem 1,*

$$mst(G) \leq \frac{n}{\delta} \left( \zeta(3) + \frac{1}{2} + \epsilon_5 \right)$$

where  $\epsilon_5 = \epsilon_5(\delta) \rightarrow 0$  as  $\delta \rightarrow \infty$ .

We prove instead

**Theorem 4** *If  $G$  is a connected graph then*

$$mst(G) \leq \frac{n}{\delta} (\zeta(3) + 1 + \epsilon_6)$$

where the  $\epsilon_6 = \epsilon_6(\delta) \rightarrow 0$  as  $\delta \rightarrow \infty$ .

We finally note that high connectivity is not necessary to obtain the result of Theorem 2. Since if  $r = o(n)$  then one can tolerate a few small cuts. For example, let  $G$  be a graph which satisfies the conditions of Theorem 2 and suppose  $r = o(n)$ . Then taking 2 disjoint copies of  $G$  and adding a single edge joining them we obtain a graph  $G'$  for which  $mst(G') \sim \frac{1}{2} + \frac{n'}{r} \zeta(3) \sim \frac{n'}{r} \zeta(3)$  where  $n' = 2n$  is the number of vertices of  $G'$ .

## 2 Proof of Theorem 3

Given a connected graph  $G = (V, E)$  with  $|V| = n$  and  $0 \leq p \leq 1$ , let  $G_p$  be the random subgraph of  $G$  with the same vertex set which contains those edges  $e$  with  $X_e \leq p$ . Let  $\kappa(G)$  denote the number of components of  $G$ . We shall first give a rather precise description of  $mst(G)$ .

**Lemma 1** [1]

*For any connected graph  $G$ ,*

$$mst(G) = \int_{p=0}^1 \mathbf{E}(\kappa(G_p)) dp - 1. \tag{3}$$

□

We substitute  $p = x/r$  in (3) to obtain

$$mst(G) = \frac{1}{r} \int_{x=0}^r \mathbf{E}(\kappa(G_{x/r})) dx - 1.$$

Now let  $C_{k,x}$  denote the total number of components in  $G_{x/r}$  with  $k$  vertices. Thus

$$mst(G) = \frac{1}{r} \int_{x=0}^r \sum_{k=1}^n \mathbf{E}(C_{k,x}) dx - 1. \tag{4}$$

### Proof of Theorem 3

In order to use (4) we need to consider three separate ranges for  $x$  and  $k$ , two of which are satisfactorily dealt with in [1]. Let  $A = (r/\omega)^{1/3}$ ,  $B = \lfloor (Ar)^{1/4} \rfloor$  so that each of  $B\alpha$ ,  $AB^2/r$  and  $A/B \rightarrow 0$  as  $r \rightarrow \infty$ . These latter conditions are needed for the analysis of the first two ranges.

**Range 1:**  $0 \leq x \leq A$  and  $1 \leq k \leq B$  – see [1].

$$\frac{1}{r} \int_{x=0}^A \sum_{k=1}^B \mathbf{E}(C_{k,x}) dx \leq (1 + o(1)) \frac{n}{r} \zeta(3).$$

**Range 2:**  $0 \leq x \leq A$  and  $k > B$  – see [1].

$$\frac{1}{r} \int_{x=0}^A \sum_{k=B}^n \mathbf{E}(C_{k,x}) dx = o(n/r).$$

**Range 3:**  $x \geq A$ .

We use a result of Karger [4]. A cut  $(S : \bar{S}) = \{(u, v) \in E : u \in S, v \notin S\}$  of  $G$  is  $\gamma$ -minimal if  $|(S : \bar{S})| \leq \gamma\lambda$ . Karger proved that the number of  $\gamma$ -minimal cuts is  $O(n^{2\gamma})$ . We can associate each component of  $G_p$  with a cut of  $G$ . Thus

$$\begin{aligned} \sum_{k=1}^n \mathbf{E}(C_{k,x}) &\leq O\left(\sum_{s=\lambda}^{\infty} n^{2s/\lambda} \left(1 - \frac{x}{r}\right)^s\right) = O\left(\sum_{s=\lambda}^{\infty} (n^{2r/\lambda} e^{-x})^{s/r}\right) \\ &= O\left(\int_{s=\lambda}^{\infty} (n^{2r/\lambda} e^{-x})^{s/r} ds\right) = O\left(\frac{rn^2 e^{-x\lambda/r}}{x - \frac{2r}{\lambda} \log n}\right), \end{aligned}$$

and

$$\frac{1}{r} \int_{x=A}^r \sum_{k=1}^n \mathbf{E}(C_{k,x}) dx = O\left(\int_{x=A}^r \frac{n^2 e^{-x\lambda/r}}{x - \frac{2r}{\lambda} \log n} dx\right) = o\left(\frac{n^2 e^{-A\lambda/r}}{r}\right) = o(n/r).$$

We complete the proof by applying Lemma 1. □

## 3 Proof of Theorem 4

We keep the definitions of  $A, B$  and Ranges 1,2, but we split Range 3 and let  $\delta = r$ .

**Range 3a:**  $x \geq A$  and  $k \leq (1 - \epsilon)r$ ,  $0 < \epsilon < 1$ , arbitrary – see [1] (here  $\epsilon = 1/2$  but the argument works for arbitrary  $\epsilon$ ).

$$\frac{1}{r} \int_{x=A}^r \sum_{k=1}^{(1-\epsilon)r} \mathbf{E}(C_{k,x}) dx = o(n/r).$$

**Range 3b:**  $x \geq A$  and  $k > (1 - \epsilon)r$ .

Clearly

$$\sum_{k=(1-\epsilon)r}^n C_{k,x} \leq \frac{n}{(1-\epsilon)r}$$

and hence

$$\frac{1}{r} \int_{x=A}^r \sum_{k=(1-\epsilon)r}^n \mathbf{E}(C_{k,x}) dx \leq \frac{n}{(1-\epsilon)r}.$$

We again complete the proof by applying Lemma 1. □

## References

- [1] A. Beveridge, A. M. Frieze and C. J. H. McDiarmid, *Minimum length spanning trees in regular graphs*, *Combinatorica* 18 (1998) 311-333.
- [2] A. M. Frieze, *On the value of a random minimum spanning tree problem*, *Discrete Applied Mathematics* 10 (1985) 47 - 56.
- [3] A. M. Frieze and C. J. H. McDiarmid, *On random minimum length spanning trees*, *Combinatorica* 9 (1989) 363 - 374.
- [4] D. R. Karger, *A Randomized Fully Polynomial Time Approximation Scheme for the All Terminal Network Reliability Problem*, Proceedings of the twenty-seventh annual ACM Symposium on Theory of Computing (1995) 11-17.
- [5] M. Penrose, *Random minimum spanning tree and percolation on the n-cube*, *Random Structures and Algorithms* 12 (1998) 63 - 82.