

Announcements

Assignments


- HW7 Due tonight!
- HW8 Due Tues 3/28, 10 pm
- P4: MDPs/RL Due Thursday 4/6, 10 pm *out!*
- Check the Final Exam Schedule for conflicts by next week 3/31

Coming up

- Midterm 2 Review – Tues. 3/28 6-8pm in GHC Rashid
- Midterm 2 – 3/30 (covers Logic, Classical Planning, MDPs, RL, Bayes)
 - 80 minutes in class, same general rules as before
 - Single function calculators or graphing calculators (no phones, iPads, etc)
 - 1 8.5"x11" handwritten pen/pencil/paper cheatsheet, not written digitally and printed

AI: Representation and Problem Solving

Bayes Nets




Instructor: Stephanie Rosenthal

Slide credits: CMU AI and <http://ai.berkeley.edu>

Omega Pizzeria!

What is the probability of getting a slice with:

- 1) No mushrooms
 - 2) Spinach and no mushrooms
 - 3) Spinach, when asking for slice with no mushrooms
- Mushrooms
 - Spinach
 - No spinach
 - No spinach and mushrooms
 - No spinach when asking for no mushrooms
 - No spinach when asking for mushrooms
 - Spinach when asking for mushrooms



Icons: CC, <https://openclipart.org/detail/296791/pizza-slice>

Probability Notation

$$P(A) = \frac{a_1}{a_2}$$

$$B = \{+b, -b\}$$

$$A = \{a_1, a_2, a_3\}$$

Notation and conventions in this course

$$P(B = +b, C) = \sum_{a \in \{a_1, a_2, a_3\}} P(A = a, B = +b, C)$$

$$\rightarrow P(+b, C) = \sum_{a \in \{a_1, a_2, a_3\}} P(a, +b, C) \quad \text{table}$$

■ Random variables:

- Capitalized

- Represents all potential outcomes

- e.g. C

■ Outcomes (values):

- lower case

- e.g. $+b, a_1, a_2, a_3$

■ Variables for values:

- lower case

- E.g. a

$$P(A, +b, C)$$

	$+b$	C	
a_1	$+b$	$+c$	✓
a_1	$+b$	$-c$	✓
a_1	$-b$	$+c$	✓
a_1	$-b$	$-c$	✓
a_2	$+b$	$+c$	✓
a_2	$+b$	$-c$	✓
a_3	$+b$	$+c$	✓

	b	c	$P(a, b, c)$
a_1	$+b$	$+c$	✓
a_1	$+b$	$-c$	✓
a_1	$-b$	$+c$	✓
a_1	$-b$	$-c$	✓
a_2	$+b$	$+c$	✓
a_2	$+b$	$-c$	✓
a_3	$+b$	$+c$	✓

Probability Notation

Notation and conventions in this course

$$P(B = +b, C) = \sum_{a \in \{a_1, a_2, a_3\}} P(A = a, B = +b, C)$$

$$P(+b, C) = \sum_{a \in \{a_1, a_2, a_3\}} P(a, +b, C)$$

Discrete Probability Distributions

Partitions

For each random variable

- Discrete outcomes
- Disjoint outcomes
- Accounts for entire event space
- Not always binary




Discrete Random Variables
(and their domains)

$$A \in \{a_1, a_2, a_3\} \leftarrow$$

$$B \in \{+b, -b\}$$

$$C \in \{+c, -c\}$$

Event space



Discrete Probability Distributions



Discrete Random Variables
(and their domains)

$$A \in \{a_1, a_2, a_3\}$$

$$B \in \{+b, -b\}$$

$$C \in \{+c, -c\}$$


Marginal distribution



Discrete Probability Distributions

Joint distribution

$P(A, B, C)$



Discrete Random Variables
(and their domains)

$A \in \{a_1, a_2, a_3\}$

$B \in \{+b, -b\}$

$C \in \{+c, -c\}$

12 rows

Discrete Probability Distributions

Joint distribution

$P(M, S, R)$

Discrete Random Variables
(and their domains)

$M \in \{m_1, m_2\}$ 2

$S \in \{s_1, s_2\}$ $\times 2$

$R \in \{r_1, r_2\}$ $\times 2$




M	S	R	$P(M, S, R)$
tm	ts	tr	2/20
tm	ts	-r	2/20
tm	-s	tr	2/20
tm	-s	-r	2/20
-m	ts	tr	1/20
-m	ts	-r	5/20
-m	-s	tr	1/20
-m	-s	-r	5/20

Icons: CC, <https://openclipart.org/detail/296791/pizza-slice>

Discrete Probability Distributions

Conditional distribution

$$P(M, S | r_2)$$



Discrete Random Variables
(and their domains)

$$M \in \{m_1, m_2\}$$

$$S \in \{s_1, s_2\}$$


$$R \in \{r_1, r_2\}$$

M	S	$P(M, S r_2)$
$+m$	$+s$	$2/6$
$+m$	$-s$	$2/6$
$-m$	$+s$	$1/6$
$-m$	$-s$	$1/6$

Discrete Probability Distributions

Conditional distribution

$$P(M, S | r_2)$$



Discrete Random Variables
(and their domains)

$$M \in \{m_1, m_2\}$$


$$S \in \{s_1, s_2\}$$

$$R \in \{r_1, r_2\}$$

Discrete Probability Distributions

Conditional distribution

$$P(\underline{A, B} \mid +c)$$



Discrete Random Variables
(and their domains)

$$A \in \{a_1, a_2, a_3\}$$


$$B \in \{+b, -b\}$$

$$C \in \{+c, -c\}$$

Discrete Probability Distributions

Conditional distribution

$$P(A, B \mid -c)$$



Discrete Random Variables
(and their domains)

$$A \in \{a_1, a_2, a_3\}$$

$$B \in \{+b, -b\}$$


$$C \in \{+c, -c\}$$

Poll 1

Which of the following probability tables sum to one?

Select all that apply.

- i. $P(\underline{A} \mid \underline{b})$
- ii. $P(\underline{A}, \underline{b}, \underline{C}) \leftarrow P(b)$
- iii. $P(\underline{A}, \underline{C} \mid \underline{b})$
- iv. $P(\underline{a}, \underline{c} \mid \underline{b})$
- v. $P(\underline{a} \mid \underline{B}, \underline{C})$
- vi. $P(\underline{c} \mid \underline{A})$



	+b
	-b

Poll 1

Which of the following probability tables sum to one?

Select all that apply.

i. $P(A | b)$

ii. $P(A, b, C)$

iii. $P(A, C | b)$

iv. $P(a, b | c)$

v. $P(a | B, C)$

vi. $P(c | A)$

Poll 2

How many valid equations can we compose using:

$P(x), P(y), P(x,y), P(x|y), P(y|x)$ and $=, \times, \div$

Bayes Rule

First one:

$$\left. \begin{aligned} P(x|y) &= P(x,y)/P(y) \\ P(y|x) &= P(x,y)/P(x) \end{aligned} \right\} \begin{array}{l} \text{def} \\ \text{cond} \\ \text{prob} \end{array}$$

$$\left. \begin{aligned} P(x|y)P(y) &= P(x,y) \\ P(y|x)P(x) &= P(x,y) \end{aligned} \right\} \text{Product rule}$$

$$\underline{P(x|y)P(y) = P(y|x)P(x)} \quad \text{Chain rule}$$

$$\frac{P(x|y)P(y)}{P(x,y)} = 1$$

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$

At most one use per probability term

e.g. Not $P(x) = P(x)$

Must be different

e.g. Cannot also use

$$P(x,y)/P(y) = P(x|y)$$

A) 2

B) 4

C) 7

D) >7

E) Other

Poll 2

Also (less meaningful): $P(y) = P(y|x)P(x) / P(x|y)$ ^(x2)
 $P(y|x) / P(x|y) = P(y) / P(x)$ ^(x2)

How many valid equations can we compose using:

$P(x)$, $P(y)$, $P(x, y)$, $P(x|y)$, $P(y|x)$ and $=$, \times , \div

First one: $P(x|y) = P(x, y) / P(y)$

$$P(y|x) = P(x, y) / P(x)$$

$$P(x, y) = P(y|x) P(x)$$

$$P(x, y) = P(x|y) P(y)$$

$$P(y|x)P(x) = P(x|y)P(y)$$

$$P(y|x) = P(x|y)P(y) / P(x)$$

$$P(x|y) = P(y|x)P(x) / P(y)$$

A) 2

B) 4

C) 7

D) >7

E) Other

At most one use per probability term

e.g. Not $P(x) = P(x)$

Must be different

e.g. Cannot also use

$$P(x, y) / P(y) = P(x|y)$$

Probability Tools Summary

Our toolbox

1. Definition of conditional probability

$$P(A|B) = \frac{P(A, B)}{P(B)}$$

2. Product Rule

$$P(A, B) = P(A|B)P(B)$$

3. Bayes' theorem

$$P(c) P(B|c) P(A|B, c) P(B|A) = \frac{P(A|B)P(B)}{P(A)}$$

4. Chain Rule

$$P(\underline{X_1, \dots, X_N}) = \prod_{n=1}^N P(X_n | X_1, \dots, X_{n-1})$$


C, B, A

$$P(A)P(B|A)P(C|A, B)$$

Answer Any Query from Joint Distribution

What is the probability of getting a slice with:

- 1) No mushrooms
 - 2) Spinach and no mushrooms
 - 3) Spinach, when asking for slice with no mushrooms
- Mushrooms
 - Spinach
 - No spinach
 - No spinach and mushrooms
 - No spinach when asking for no mushrooms
 - No spinach when asking for mushrooms
 - Spinach when asking for mushrooms



Icons: CC, <https://openclipart.org/detail/296791/pizza-slice>

Answer Any Query from Joint Distribution

You can answer all of these questions:

$P(M)$	
m_1	12/20
m_2	

$P(S)$	
s_1	
s_2	

$P(M, S)$		
m_1	s_1	
m_1	s_2	6/20
m_2	s_1	
m_2	s_2	

$P(M s_1)$	
m_1	
m_2	

$P(M s_2)$	
m_1	
m_2	

$P(S m_1)$	
s_1	
s_2	6/12

$P(S m_2)$	
s_1	
s_2	

Answer Any Query from Joint Distribution

P(Weather)?

W	P(W)
S	.65
r	1-.65

$\sum_{\{S, w\}} \sum_{Temp} P(S, T, W)$

P(Weather | winter)?

W	P(w winter)
S	.5
r	.5

$\frac{\sum_{T} P(winter, T, w)}{P(winter)}$

$P(winter) = .5$

P(Weather | winter, hot)?

Season	Temp	Weather	P(S, T, W)
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

Answer Any Query from Joint Distribution

P(Weather)?

Season	Temp	Weather	P(S, T, W)
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

Answer Any Query from Joint Distribution

$P(\text{Weather} \mid \text{winter})?$

Season	Temp	Weather	$P(S, T, W)$
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

Answer Any Query from Joint Distribution

$P(\text{Weather} \mid \text{winter, hot})?$

Season	Temp	Weather	$P(S, T, W)$
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

Additional Probability Tools

- Marginalization (law of total probability) (summing out)

$$\rightarrow P(A) = \sum_b \sum_c P(A, b, c)$$

- Normalization

$$\underline{P(B | a)} = \frac{P(a, B)}{P(a)}$$

$$\underline{P(B | a)} \propto \underline{P(a, B)}$$

$$\underline{P(B | a)} = \frac{1}{\underline{z}} P(a, B)$$

$$\alpha = \frac{1}{z}$$


$$\underline{z} = \underline{P(a)} = \sum_b P(a, b)$$

$$\frac{P(a, b)}{P(a)}$$

Answer Any Query from Joint Distribution

Joint distributions are the best!

Joint





query

Query

evidence

$$\underline{P(q_1, q_2 \mid e_1, e_2, e_3)}$$

Answer Any Query from Joint Distribution

Two tools to go from joint to query

1. Definition of conditional probability

$$P(A|B) = \frac{P(A, B)}{P(B)}$$

2. Law of total probability (marginalization, summing out)

$$P(A) = \sum_b P(A, b)$$

$$P(Y | U, V) = \sum_x \sum_z P(x, Y, z | U, V)$$

Answer Any Query from Joint Distribution

Two tools to go from joint to query

Joint: $P(H_1, H_2, Q, E)$

Query: $P(Q | e)$

1. Definition of conditional probability

$$P(Q|e) = \frac{P(Q, e)}{\underline{P(e)}}$$

2. Law of total probability (marginalization, summing out)

$$P(Q, e) = \sum_{h_1} \sum_{h_2} P(h_1, h_2, Q, e)$$

hidden

$$\underline{P(e)} = \sum_q \sum_{h_1} \sum_{h_2} P(h_1, h_2, q, e)$$

Answer Any Query from Joint Distribution

$P(\text{Weather})?$

$P(\text{Weather} \mid \text{winter})?$

$P(\text{Weather} \mid \text{winter, hot})?$

Season	Temp	Weather	$P(S, T, W)$
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20


Answer Any Query from Joint Distribution

Joint distributions are the best!

Problems with joints

- We aren't given the joint table
 - Usually some set of conditional probability tables

Joint






Query


$$P(a | e)$$

Build Joint Distribution Using Chain Rule

Conditional Probability Tables
and Chain Rule



easier to compute



Joint

$$P(A) P(B|A) P(C|A, B) P(D|A, B, C) P(E|A, B, C, D)$$

Query
 $P(a | e)$

Build Joint Distribution Using Chain Rule

Two tools to construct joint distribution

1. Product rule

$$P(A, B) = P(A | B)P(B)$$
$$P(A, B) = P(B | A)P(A)$$

2. Chain rule

$$P(X_1, X_2, \dots, X_n) = \prod_i P(X_i | X_1, \dots, X_{i-1})$$

$$P(A, B, C) = P(A)P(B | A)P(C | A, B) \quad \text{for ordering A, B, C}$$

$$P(A, B, C) = P(A)P(C | A)P(B | A, C) \quad \text{for ordering A, C, B}$$


$$P(A, B, C) = P(C)P(B | C)P(A | C, B) \quad \text{for ordering C, B, A}$$

Build Joint Distribution Using Chain Rule

Binary random variables

- Fire
- Smoke
- Alarm

$$P(F, S, A)$$



$$\textcircled{1} P(F)P(S|F)P(A|F,S)$$


$$\textcircled{2} P(S)P(A|S)P(F|A,S)$$

$$6 = 3 \times 2$$

Poll 3

Variables

- B: Burglary
- A: Alarm goes off
- M: Mary calls
- J: John calls
- E: Earthquake!




How many different ways can we write the chain rule?

- A. 1
- B. 5
- C. 5 choose 5
- D. 5!
- E. 5^5


$$5 \times 4 \times 3 \times 2 \times 1$$

Build Joint Distribution Using Chain Rule

Conditional Probability Tables
and Chain Rule



Joint



Query

$$P(a | e)$$

$$P(A) P(B|A) P(C|A, B) P(D|A, B, C) P(E|A, B, C, D)$$

Answer Any Query from Condition Probability Tables

Process to go from (specific) conditional probability tables to query

1. Construct the joint distribution
 1. Product Rule or Chain Rule
2. Answer query from joint
 1. Definition of conditional probability
 2. Law of total probability (marginalization, summing out)

Answer Any Query from Condition Probability Tables

Bayes' rule as an example

Given: $P(E|Q)$, $P(Q)$ Query: $P(Q | e)$

1. Construct the **joint** distribution

1. Product Rule or Chain Rule

$$P(E, Q) = P(E|Q)P(Q)$$

2. Answer **query** from **joint**

1. Definition of conditional probability

$$P(Q | e) = \frac{P(e, Q)}{P(e)}$$

2. Law of total probability (marginalization, summing out)


$$P(Q | e) = \frac{P(e, Q)}{\sum_q P(e, q)}$$

Bayesian Networks

One node per random variable

DAG

One CPT per node: $P(\text{node} \mid \text{Parents}(\text{node}))$




$$P(A, B, C, D) = \underbrace{P(A)} \underbrace{P(B|A)} P(C|A, B) P(D|A, B, C)$$

Encode joint distributions as product of conditional distributions on each variable

$$\rightarrow P(X_1, \dots, X_N) = \prod_i P(X_i \mid \text{Parents}(X_i))$$

Bayes net




tail
head


Build Bayes Net Using Chain Rule

Binary random variables

- Fire
- Smoke
- Alarm





$$P(F) P(S|F) P(A|S, F)$$



Question

Variables



- B: Burglary
- A: Alarm goes off
- M: Mary calls
- J: John calls
- E: Earthquake!




Given the Bayes net, write the joint distribution?

Answer Any Query from Bayes Net

Bayes Net and
Conditional
Probability Tables



Joint




Query


$$P(a | e)$$

Answer Any Query from Condition Probability Tables

Conditional Probability Tables
and Chain Rule



Joint




Query

$$P(a | e)$$

$$P(A) P(B|A) P(C|A, B) P(D|A, B, C) P(E|A, B, C, D)$$

Answer Any Query from Condition Probability Tables

Conditional Probability Tables and Chain Rule



$$P(A) \quad P(B|A) \quad P(C|A, B) \quad P(D|A, B, C) \quad P(E|A, B, C, D)$$


Problems

- Huge
 - n variables with d values
 - d^n entries
- We aren't given the right tables

Do We Need the Full Chain Rule?


Binary random variables

- Fire
- Smoke
- Alarm




Answer Any Query from Condition Probability Tables

Conditional Probability Tables
and Chain Rule



Joint





Query

$$P(a | e)$$

$$P(A) P(B|A) P(C|A, B) P(D|A, B, C) P(E|A, B, C, D)$$

Answer Any Query from Condition Probability Tables


Bayes Net



$P(A)$ $P(B|A)$ $P(C|A)$ $P(D|C)$ $P(E|C)$

$$P(X_1, \dots, X_N) = \prod_i P(X_i | \text{Parents}(X_i))$$

Joint





Query

$P(a | e)$