

CARNEGIE MELLON UNIVERSITY
DEPARTMENT OF COMPUTER SCIENCE
15-826 MULTIMEDIA DATABASES AND DATA MINING
C. FALOUTSOS, FALL 2024

Homework 3

Due: pdf, on canvas, at 2:00pm, on 11/01/2024

VERY IMPORTANT:

- Upload **e-copy** of your answers, on canvas.

Reminders:

- *Plagiarism*: Homework is to be completed *individually*.
- *Typeset* your answers. Illegible handwriting may get zero points.
- *Late homeworks*: Follow the published policy

For your information:

- Explanations are *optional*, and will only be used to for partial credit, if the main answer is off.
- Graded out of **100** points; **5** questions total
- Rough time estimate: *1-2 hours ($\approx 10-20$ minutes per question)*

Revision : 2024/10/04 02:39

Question	Points	Score
R-trees	10	
Fractals - warm-up	20	
Fractals - T/F	20	
Correlation integrals	25	
Power laws	25	
Total:	100	

Question 1: R-trees [10 points]

A parent node in an R-tree of $d=4$ dimensions, has an MBR with sizes $x_1=0.1$, $x_2=0.2$, $x_3=0.3$ and $x_4=0.4$.

Remember: Explanations are *optional*, and will only be used to for partial credit, if the main answer is off.

- (a) **[10 points]** What is the probability it will be retrieved by a point query? (Assume uniform distribution of queries; assume the address space is the unit cube).

(a) _____

.....
.....
.....
.....

Question 2: Fractals - warm-up..... [20 points]

Remember: Explanations are *optional*, and will only be used to for partial credit, if the main answer is off.

- (a) **[10 points]** Consider 1M points on the Koch snowflake (see course lectures, slides #13-17, or wikipedia). Suppose that point P has $n_1=10$ neighbors within $r_1=0.001$ unit of distance. How many neighbors do you expect it to have, when we triple the radius, that is, how many neighbors within $r_2=3*r_1$ units of distance?

(a) _____

.....

- (b) **[10 points]** Same as above, but with 1M points on the Cantor Dust (see course lectures, slides #19-20): Point Q has $m_1=10$ neighbors within $r_1=0.001$. How many neighbors do you expect it to have, within $r_2=3*r_1$ units of distance?

(b) _____

.....

Question 3: Fractals - T/F [20 points]

Mark your response (Yes or No), for the following statements. No need for explanations. As in the class, by 'fractal dimension' we mean the 'correlation fractal dimension'.

Is it possible that a cloud of points in $E(\geq 1)$ dimensions may have:

- (a) **[2 points]** fractal dimension equal to E ?
 Yes No
- (b) **[2 points]** fractal dimension equal to $E - 1$?
 Yes No
- (c) **[2 points]** fractal dimension equal to $E/2$?
 Yes No
- (d) **[2 points]** fractal dimension equal to $E + 1$?
 Yes No
- (e) **[2 points]** fractal dimension = 1, *without* the dataset being points from a straight line?
 Yes No
- (f) **[2 points]** fractal dimension < 1 ?
 Yes No
- (g) **[2 points]** negative fractal dimension?
 Yes No
- (h) **[2 points]** fractal dimension equal to zero?
 Yes No
- (i) **[2 points]** fractal dimension = 0.5 (a rational number)?
 Yes No
- (j) **[2 points]** *several, different* fractal dimensions, in different ranges of scales?
 Yes No

Question 4: Correlation integrals [25 points]

Consider some clouds of points, some of which are shown in Figure 1. We want to describe their correlation integrals. No need for explanations.




Figure 1: Clouds of points: for 3D-STICKS we show a few 'sticks'; for CANTOR, we show the result after the first 2 iterations of eliminating the *middle third*

Each cloud has N points (say, $N=100,000$ - the exact value is not needed). The descriptions of the clouds follow. For each cloud, list the slopes (left-to-right). Figure 2 gives reminders of how correlation integrals look like.




Figure 2: Illustrations of some correlation integrals and their slopes

(a) [1 point] LINE: points uniformly distributed on the major diagonal of the unit square (zero thickness)

(a) _____

(b) [3 points] CIRCLE: points uniformly distributed along the periphery of the unit circle (zero thickness)

(b) _____

(c) [3 points] DISC: points uniformly distributed in the unit circle

(c) _____

- (d) **[6 points]** RING: points uniformly distributed inside a ring of radius=1 and width $w=0.01$ (that is, RING is like 'CIRCLE', but with non-zero thickness).

(d) _____

- (e) **[6 points]** 3D-STICKS: points on 1,000 line segments; each segment is vertical, with height 10^{-4} (and obviously, zero width); the segment-centers are uniformly distributed in the unit **cube**.

(e) _____

- (f) **[6 points]** CANTOR: points on the (zero width) diagonal line of the unit square, after eliminating the middle-third, recursively. In figure 1, we show the first two iterations of the above procedure.

(f) _____

Question 5: Power laws.....[25 points]

We are told that some country has $N=10^7$ citizens, and that their income X follows a power law:

$$P(X \geq x) = x^{-1} \quad x \geq X_{min} = 1 \quad (1)$$

That is, the CCDF (complementary cumulative distribution function) is a power law with exponent -1 , and the lowest income X_{min} is 1.

Remember: Explanations are *optional*, and will only be used to for partial credit, if the main answer is off.

- (a) [5 points] What is median income?

(a) _____

.....

- (b) [10 points] What is your estimate for the income X_{max} of the richest citizen?

(b) _____

.....

- (c) [5 points] What is your estimate for the income X_2 of the second richest person?

(c) _____

.....

- (d) [5 points] From the lectures, we know that the pdf (probability density function) of the income $p(x) \propto x^\beta$, is also a power law. What is the value of the exponent β ?

(d) _____

.....

