

Machine Learning


10-701/15-781, Spring 2010

Support Vector Machines

Eric Xing

Lecture 17, March 15, 2010

Reading: Chap. 6&7, C.B book, and listed papers




© Eric Xing @ CMU, 2006-2010

1

What is a good Decision Boundary?

- Consider a binary classification task with $y = \pm 1$ labels (not 0/1 as before).
- When the training examples are linearly separable, we can set the parameters of a linear classifier so that all the training examples are classified correctly
- Many decision boundaries!
 - Generative classifiers
 - Logistic regressions ...
- Are all decision boundaries equally good?



© Eric Xing @ CMU, 2006-2010

2



What is a good Decision Boundary?



© Eric Xing @ CMU, 2006-2010

3

Not All Decision Boundaries Are Equal!



- Why we may have such boundaries?
 - Irregular distribution
 - Imbalanced training sizes
 - outliers

© Eric Xing @ CMU, 2006-2010

4


Classification and Margin



- Parameterizing decision boundary

- Let w denote a vector orthogonal to the decision boundary, and b denote a scalar "offset" term, then we can write the decision boundary as:

$$w^T x + b = 0$$



© Eric Xing @ CMU, 2006-2010

5


Classification and Margin



- Parameterizing decision boundary

- Let w denote a vector orthogonal to the decision boundary, and b denote a scalar "offset" term, then we can write the decision boundary as:

$$w^T x + b = 0$$



- Margin

$$w^T x_i + b > +c \quad \text{for all } x_i \text{ in class 2}$$

$$w^T x_i + b < -c \quad \text{for all } x_i \text{ in class 1}$$

Or more compactly:

$$(w^T x_i + b) y_i > c$$

The margin between any two points

$$m = d^- + d^+ =$$

© Eric Xing @ CMU, 2006-2010

6

Maximum Margin Classification



- The "minimum" permissible margin is:

$$m = \frac{w^T}{\|w\|} (x_{i^*} - x_{j^*}) = \frac{2c}{\|w\|}$$

- Here is our Maximum Margin Classification problem:

$$\begin{aligned} \max_w \quad & \frac{2c}{\|w\|} \\ \text{s.t.} \quad & y_i(w^T x_i + b) \geq c, \quad \forall i \end{aligned}$$

© Eric Xing @ CMU, 2006-2010

7

Maximum Margin Classification, con'd.



- The optimization problem:

$$\begin{aligned} \max_{w,b} \quad & \frac{c}{\|w\|} \\ \text{s.t.} \quad & y_i(w^T x_i + b) \geq c, \quad \forall i \end{aligned}$$

- But note that the magnitude of c merely scales w and b , and does not change the classification boundary at all! (why?)
- So we instead work on this cleaner problem:

$$\begin{aligned} \max_{w,b} \quad & \frac{1}{\|w\|} \\ \text{s.t.} \quad & y_i(w^T x_i + b) \geq 1, \quad \forall i \end{aligned}$$

- The solution to this leads to the famous **Support Vector Machines** - -- believed by many to be the best "off-the-shelf" supervised learning algorithm

© Eric Xing @ CMU, 2006-2010

8


Support vector machine



- A convex quadratic programming problem with linear constraints:

$$\begin{aligned} \max_{w,b} \quad & \frac{1}{\|w\|} \\ \text{s.t.} \quad & y_i(w^T x_i + b) \geq 1, \quad \forall i \end{aligned}$$

- The attained margin is now given by $\frac{1}{\|w\|}$
- Only a few of the classification constraints are relevant → **support vectors**



- Constrained optimization
 - We can directly solve this using commercial quadratic programming (QP) code
 - But we want to take a more careful investigation of Lagrange duality, and the solution of the above in its dual form.
- deeper insight: support vectors, kernels ...
- more efficient algorithm

© Eric Xing @ CMU, 2006-2010

9

Digression to Lagrangian Duality



- The Primal Problem

$$\begin{aligned} \text{Primal:} \quad & \min_w \quad f(w) \\ & \text{s.t.} \quad g_i(w) \leq 0, \quad i=1, \dots, k \\ & \quad \quad h_i(w) = 0, \quad i=1, \dots, l \end{aligned}$$

The generalized Lagrangian:

$$\mathcal{L}(w, \alpha, \beta) = f(w) + \sum_{i=1}^k \alpha_i g_i(w) + \sum_{i=1}^l \beta_i h_i(w)$$

the α 's ($\alpha_i \geq 0$) and β 's are called the Lagrangian multipliers

Lemma:

$$\max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta) = \begin{cases} f(w) & \text{if } w \text{ satisfies primal constraints} \\ \infty & \text{o/w} \end{cases}$$

A re-written Primal:

$$\min_w \max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta)$$

© Eric Xing @ CMU, 2006-2010

10

Lagrangian Duality, cont.



- Recall the Primal Problem:

$$\min_w \max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta)$$

- The Dual Problem:

$$\max_{\alpha, \beta, \alpha_i \geq 0} \min_w \mathcal{L}(w, \alpha, \beta)$$

- Theorem (weak duality):**

$$d^* = \max_{\alpha, \beta, \alpha_i \geq 0} \min_w \mathcal{L}(w, \alpha, \beta) \leq \min_w \max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta) = p^*$$

- Theorem (strong duality):**

Iff there exist a saddle point of $\mathcal{L}(w, \alpha, \beta)$, we have

$$d^* = p^*$$

© Eric Xing @ CMU, 2006-2010


11

A sketch of strong and weak duality



- Now, ignoring $h(x)$ for simplicity, let's look at what's happening graphically in the duality theorems.

$$d^* = \max_{\alpha_i \geq 0} \min_w f(w) + \alpha^T g(w) \leq \min_w \max_{\alpha_i \geq 0} f(w) + \alpha^T g(w) = p^*$$



© Eric Xing @ CMU, 2006-2010


12

A sketch of strong and weak duality



- Now, ignoring $h(x)$ for simplicity, let's look at what's happening graphically in the duality theorems.

$$d^* = \max_{\alpha_i \geq 0} \min_w f(w) + \alpha^T g(w) \leq \min_w \max_{\alpha_i \geq 0} f(w) + \alpha^T g(w) = p^*$$



© Eric Xing @ CMU, 2006-2010


13

A sketch of strong and weak duality



- Now, ignoring $h(x)$ for simplicity, let's look at what's happening graphically in the duality theorems.

$$d^* = \max_{\alpha_i \geq 0} \min_w f(w) + \alpha^T g(w) \leq \min_w \max_{\alpha_i \geq 0} f(w) + \alpha^T g(w) = p^*$$



© Eric Xing @ CMU, 2006-2010

14



The KKT conditions

- If there exists some saddle point of \mathcal{L} , then the saddle point satisfies the following "Karush-Kuhn-Tucker" (KKT) conditions:

$$\frac{\partial}{\partial w_i} \mathcal{L}(w, \alpha, \beta) = 0, \quad i = 1, \dots, k$$

$$\frac{\partial}{\partial \beta_i} \mathcal{L}(w, \alpha, \beta) = 0, \quad i = 1, \dots, l$$

$$a_i g_i(w) = 0, \quad i = 1, \dots, m$$

$$g_i(w) \leq 0, \quad i = 1, \dots, m$$

$$\alpha_i \geq 0, \quad i = 1, \dots, m$$

Complementary slackness

Primal feasibility

Dual feasibility

- Theorem:** If w^* , α^* and β^* satisfy the KKT condition, then it is also a solution to the primal and the dual problems.



Solving optimal margin classifier

- Recall our opt problem:

$$\begin{aligned} & \max_{w,b} \frac{1}{\|w\|} \\ \text{s.t.} & \quad y_i(w^T x_i + b) \geq 1, \quad \forall i \end{aligned}$$

- This is equivalent to

$$\begin{aligned} & \min_{w,b} \frac{1}{2} w^T w \\ \text{s.t.} & \quad 1 - y_i(w^T x_i + b) \leq 0, \quad \forall i \end{aligned} \quad (*)$$

- Write the Lagrangian:

$$\mathcal{L}(w, b, \alpha) = \frac{1}{2} w^T w - \sum_{i=1}^m \alpha_i [y_i(w^T x_i + b) - 1]$$

- Recall that (*) can be reformulated as $\min_{w,b} \max_{\alpha_i \geq 0} \mathcal{L}(w, b, \alpha)$
Now we solve its **dual problem**: $\max_{\alpha_i \geq 0} \min_{w,b} \mathcal{L}(w, b, \alpha)$

The Dual Problem



$$\max_{\alpha_i \geq 0} \min_{w, b} \mathcal{L}(w, b, \alpha)$$

- We minimize \mathcal{L} with respect to w and b first:

$$\nabla_w \mathcal{L}(w, b, \alpha) = w - \sum_{i=1}^m \alpha_i y_i x_i = 0, \quad (*)$$

$$\nabla_b \mathcal{L}(w, b, \alpha) = \sum_{i=1}^m \alpha_i y_i = 0, \quad (**)$$

Note that (*) implies: $w = \sum_{i=1}^m \alpha_i y_i x_i \quad (***)$

- Plus (***) back to \mathcal{L} , and using (**), we have:

$$\mathcal{L}(w, b, \alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

© Eric Xing @ CMU, 2006-2010

17

The Dual problem, cont.



- Now we have the following dual opt problem:

$$\max_{\alpha} \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{s.t. } \alpha_i \geq 0, \quad i = 1, \dots, k$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- This is, (again,) a **quadratic programming** problem.

- A global maximum of α_i can always be found.
- But what's the big deal??
- Note two things:

1. w can be recovered by $w = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i$ See next ...

2. The "kernel" $\mathbf{x}_i^T \mathbf{x}_j$ More later ...

© Eric Xing @ CMU, 2006-2010


18

Support vectors



- Note the KKT condition --- only a few α_i 's can be nonzero!!

$$\alpha_i g_i(w) = 0, \quad i = 1, \dots, m$$



© Eric Xing @ CMU, 2006-2010

19

Support vector machines



- Once we have the Lagrange multipliers $\{\alpha_i\}$, we can reconstruct the parameter vector w as a weighted combination of the training examples:

$$w = \sum_{i \in SV} \alpha_i y_i \mathbf{x}_i$$

- For testing with a new data z

- Compute

$$w^T z + b = \sum_{i \in SV} \alpha_i y_i (\mathbf{x}_i^T z) + b$$

and classify z as class 1 if the sum is positive, and class 2 otherwise

- Note: w need not be formed explicitly

© Eric Xing @ CMU, 2006-2010

20

Interpretation of support vector machines




- The optimal w is a linear combination of a small number of data points. This “sparse” representation can be viewed as data compression as in the construction of kNN classifier
- To compute the weights $\{\alpha_i\}$, and to use support vector machines we need to specify only the inner products (or kernel) between the examples $x_i^T x_j$
- We make decisions by comparing each new example z with only the support vectors:

$$y^* = \text{sign} \left(\sum_{i \in SV} \alpha_i y_i (x_i^T z) + b \right)$$

© Eric Xing @ CMU, 2006-2010

21

Non-linearly Separable Problems



- We allow “error” ξ_i in classification; it is based on the output of the discriminant function $w^T x + b$
- ξ_i approximates the number of misclassified samples

© Eric Xing @ CMU, 2006-2010

22

Soft Margin Hyperplane



- Now we have a slightly different opt problem:

$$\begin{aligned} \min_{w,b} \quad & \frac{1}{2} w^T w + C \sum_{i=1}^m \xi_i \\ \text{s.t.} \quad & y_i(w^T x_i + b) \geq 1 - \xi_i, \quad \forall i \\ & \xi_i \geq 0, \quad \forall i \end{aligned}$$

- ξ_i are “slack variables” in optimization
- Note that $\xi_i=0$ if there is no error for \mathbf{x}_i
- ξ_i is an upper bound of the number of errors
- C : tradeoff parameter between error and margin

© Eric Xing @ CMU, 2006-2010

23

The Optimization Problem



- The dual of this new constrained optimization problem is

$$\begin{aligned} \max_{\alpha} \quad & \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j) \\ \text{s.t.} \quad & 0 \leq \alpha_i \leq C, \quad i=1, \dots, m \\ & \sum_{i=1}^m \alpha_i y_i = 0. \end{aligned}$$

- This is very similar to the optimization problem in the linear separable case, except that there is an upper bound C on α_i now
- Once again, a QP solver can be used to find α_i

© Eric Xing @ CMU, 2006-2010

24

The SMO algorithm




- Consider solving the **unconstrained** opt problem:

$$\max_{\alpha} W(\alpha_1, \alpha_2, \dots, \alpha_m)$$

- We've already see three opt algorithms!
 - ?
 - ?
 - ?
- Coordinate ascend:

Coordinate ascend



Sequential minimal optimization



- Constrained optimization:

$$\max_{\alpha} \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{s.t. } 0 \leq \alpha_i \leq C, \quad i = 1, \dots, m$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- Question: can we do coordinate along one direction at a time (i.e., hold all $\alpha_{[-i]}$ fixed, and update α_i ?)

The SMO algorithm



Repeat till convergence

1. Select some pair α_i and α_j to update next (using a heuristic that tries to pick the two that will allow us to make the biggest progress towards the global maximum).
2. Re-optimize $J(\alpha)$ with respect to α_i and α_j , while holding all the other α_k 's ($k \neq i, j$) fixed.

Will this procedure converge?

Convergence of SMO



$$\max_{\alpha} \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{KKT:} \quad \text{s.t.} \quad 0 \leq \alpha_i \leq C, \quad i=1, \dots, k$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- Let's hold $\alpha_3, \dots, \alpha_m$ fixed and reopt J w.r.t. α_1 and α_2

Convergence of SMO



- The constraints:

$$\alpha_1 y_1 + \alpha_2 y_2 = \xi$$


$$0 \leq \alpha_1 \leq C$$

$$0 \leq \alpha_2 \leq C$$

- The objective:

$$\mathcal{J}(\alpha_1, \alpha_2, \dots, \alpha_m) = \mathcal{J}((\xi - \alpha_2 y_2) y_1, \alpha_2, \dots, \alpha_m)$$

- Constrained opt:




Cross-validation error of SVM



- The leave-one-out cross-validation error does not depend on the dimensionality of the feature space but only on the # of support vectors!

$$\text{Leave-one-out CV error} = \frac{\# \text{ support vectors}}{\# \text{ of training examples}}$$



© Eric Xing @ CMU, 2006-2010

31

Summary



- Max-margin decision boundary
- Constrained convex optimization
 - Duality
 - The KKT conditions and the support vectors
 - Non-separable case and slack variables
 - The SMO algorithm

© Eric Xing @ CMU, 2006-2010

32